

New Ostrowski type inequalities for harmonically convex functions

Tuncay Koroglu

Karadeniz Technical University, Department of Mathematics, Trabzon, Turkey

Received: 11 December 2017, Accepted: 15 December 2017

Published online: 7 March 2018.

Abstract: In this work, we pose a new equality for differentiable functions. By using this equality, we have some new Ostrowski type inequalities and some error estimates for the midpoint formula for functions whose derivatives in absolute values at certain powers are harmonically convex.

Keywords: Ostrowski type inequalities, midpoint type inequalities, harmonically convex function.

1 Introduction

The following result is known in the literature as Ostrowski's inequality [7];

Theorem 1. [2,8]. *Let $f : [a,b] \rightarrow \mathbb{R}$ be a differentiable mapping on (a,b) with the property that $|f'(t)| \leq M$ for all $t \in (a,b)$. Then*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq (b-a)M \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] \quad (1)$$

for all $x \in [a,b]$.

The constant $\frac{1}{4}$ is the best possible it means that it cannot be replaced by a smaller constant.

The inequality (1) can be expressed in the following form:

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{M}{b-a} \left[\frac{(x-a)^2 + (x-b)^2}{2} \right]. \quad (2)$$

For some results which generalize, improve and extend the inequalities (1) and (2) we refer the reader to the recent papers (see [1,2,4,5,7,8,9]).

In [3], İşcan gave definition of harmonically convex functions and Hermite-Hadamard type inequality for harmonically convex functions as follows:

Definition 1. [3]. *Let $I \subset \mathbb{R} \setminus \{0\}$ be a real interval. A function $f : I \rightarrow \mathbb{R}$ is said to be harmonically convex, if*

$$f\left(\frac{xy}{tx+(1-t)y}\right) \leq tf(y) + (1-t)f(x) \quad (3)$$

* Corresponding author e-mail: tkor@ktu.edu.tr

© 2018 BISKA Bilisim Technology

for all $x, y \in I$ and $t \in [0, 1]$. If the inequality in (3) is reversed, then f is said to be harmonically concave.

Theorem 2. [3]. Let $f : I \subset \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ be a harmonically convex function and $a, b \in I$ with $a < b$. If $f \in L[a, b]$ then the following inequalities holds:

$$f\left(\frac{2ab}{a+b}\right) \leq \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \leq \frac{f(a) + f(b)}{2}.$$

We will use the following special functions called beta function and hypergeometric function, respectively in the literature

$$\beta(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} = \int_0^1 t^{x-1} (1-t)^{y-1} dt, \quad x, y > 0,$$

$${}_2F_1(a, b; c; z) = \frac{1}{\beta(b, c-b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} (1-zt)^{-a} dt,$$

where $c > b > 0$, and $|z| < 1$ (see e.g. [6]).

2 New Ostrowski type inequalities

The following lemma is important to prove our main results.

Lemma 1. Let $f : I \subset \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $a, b \in I$ with $a < b$. If $f' \in L[a, b]$ then

$$f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du = ab(b-a) \int_0^1 \frac{p(t)}{(ta+(1-t)b)^2} f'\left(\frac{ab}{ta+(1-t)b}\right) dt \quad (4)$$

where

$$p(t) = \begin{cases} t, & t \in \left[0, \frac{(x-a)b}{(b-a)x}\right] \\ t-1, & t \in \left(\frac{(x-a)b}{(b-a)x}, 1\right] \end{cases}$$

for all $x \in [a, b]$.

Proof.

$$\begin{aligned} ab(b-a) \int_0^1 \frac{p(t)}{(ta+(1-t)b)^2} f'\left(\frac{ab}{ta+(1-t)b}\right) dt &= ab(b-a) \left[\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} f'\left(\frac{ab}{ta+(1-t)b}\right) dt \right. \\ &\quad \left. + \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{t-1}{(ta+(1-t)b)^2} f'\left(\frac{ab}{ta+(1-t)b}\right) dt \right] \\ &= \left[tf\left(\frac{ab}{ta+(1-t)b}\right) \Big|_0^{\frac{(x-a)b}{(b-a)x}} - \int_0^{\frac{(x-a)b}{(b-a)x}} f\left(\frac{ab}{ta+(1-t)b}\right) dt + (t-1) f\left(\frac{ab}{ta+(1-t)b}\right) \Big|_{\frac{(x-a)b}{(b-a)x}}^1 - \int_{\frac{(x-a)b}{(b-a)x}}^1 f\left(\frac{ab}{ta+(1-t)b}\right) dt \right] \\ &= \left[\frac{(x-a)b}{(b-a)x} f(x) - \frac{ab}{b-a} \int_a^x \frac{f(u)}{u^2} du + \frac{(b-x)a}{(b-a)x} f(x) - \frac{ab}{b-a} \int_x^b \frac{f(u)}{u^2} du \right] \\ &= f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du. \end{aligned}$$

This completes the proof.

Theorem 3. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $a, b \in I$ with $a < b$ and $f' \in L$. If $|f'|$ is harmonically convex on $[a, b]$, then for all $x \in [a, b]$, we have

$$\left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a) \left[\begin{array}{l} |f'(a)| (T_1(a, b, x) + T_3(a, b, x)) \\ + |f'(b)| (T_2(a, b, x) + T_4(a, b, x)) \end{array} \right] \quad (5)$$

where

$$T_1(a, b, x) = \left[\begin{array}{l} \frac{1}{2} \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{b}{3} \left(\frac{(x-a)}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right],$$

$$T_2(a, b, x) = \frac{b}{3} \left(\frac{(x-a)}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right),$$

$$T_3(a, b, x) = \left[\begin{array}{l} \frac{b^{-2}}{6} {}_2F_1 \left(2, 1; 4; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{(x-a)b^{-1}}{(b-a)x} {}_2F_1 \left(2, 1; 2; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{b}{3} \left(\frac{(x-a)}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right],$$

$$T_4(a, b, x) = \left[\begin{array}{l} \frac{b^{-2}}{6} {}_2F_1 \left(2, 2; 4; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{2} \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \frac{b}{3} \left(\frac{(x-a)}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right].$$

Proof. By using Lemma 1 and harmonically convexity of $|f'|$, we have

$$\begin{aligned} \left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| &\leq ab(b-a) \left[\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} \left| f' \left(\frac{ab}{ta+(1-t)b} \right) \right| dt \right. \\ &\quad \left. + \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^2} \left| f' \left(\frac{ab}{ta+(1-t)b} \right) \right| dt \right] \\ &\leq ab(b-a) \left[\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} [(1-t)|f'(a)| + t|f'(b)|] dt \right. \\ &\quad \left. + \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^2} [(1-t)|f'(a)| + t|f'(b)|] dt \right] \\ &\leq ab(b-a) \left[|f'(a)| \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t(1-t)}{(ta+(1-t)b)^2} dt + |f'(b)| \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^2}{(ta+(1-t)b)^2} dt \right. \\ &\quad \left. + |f'(a)| \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{(1-t)^2}{(ta+(1-t)b)^2} dt + |f'(b)| \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{t(1-t)}{(ta+(1-t)b)^2} dt \right]. \end{aligned} \quad (6)$$

If we calculate the appearing integrals with hypergeometric functions, we have

$$\begin{aligned}
\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t(1-t)}{(ta+(1-t)b)^2} dt &= \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} dt - \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^2}{(ta+(1-t)b)^2} dt \\
&= \left(\frac{(x-a)}{(b-a)x} \right)^2 \int_0^1 u \left(1 - \left[\frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right] u \right)^{-2} du \\
&\quad - b \left(\frac{(x-a)}{(b-a)x} \right)^3 \int_0^1 u^2 \left(1 - \left[\frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right] u \right)^{-2} du \\
&= \left[\begin{array}{l} \frac{1}{2} \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{b}{3} \left(\frac{(x-a)}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right] = T_1(a, b, x), \tag{7}
\end{aligned}$$

$$\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^2}{(ta+(1-t)b)^2} dt = \frac{b}{3} \left(\frac{(x-a)}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) = T_2(a, b, x), \tag{8}$$

$$\begin{aligned}
\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{(1-t)^2}{(ta+(1-t)b)^2} dt &= \int_0^1 \frac{(1-t)^2}{(ta+(1-t)b)^2} dt - \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{(1-t)^2}{(ta+(1-t)b)^2} dt \\
&= b^{-2} \int_0^1 (1-t)^2 \left(1 - t \left(1 - \frac{a}{b} \right) \right)^{-2} dt - \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{1 - 2t + t^2}{(ta+(1-t)b)^2} dt \\
&= b^{-2} \int_0^1 (1-t)^2 \left(1 - t \left(1 - \frac{a}{b} \right) \right)^{-2} dt - \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{1}{(ta+(1-t)b)^2} dt \\
&\quad + 2 \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} dt - \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^2}{(ta+(1-t)b)^2} dt \\
&= \left[\begin{array}{l} \frac{b^{-2}}{3} {}_2F_1 \left(2, 1; 4; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{(x-a)b^{-1}}{(b-a)x} {}_2F_1 \left(2, 1; 2; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{b}{3} \left(\frac{(x-a)}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right] = T_3(a, b, x), \tag{9}
\end{aligned}$$

$$\begin{aligned}
\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{t(1-t)}{(ta+(1-t)b)^2} dt &= \int_0^1 \frac{t(1-t)}{(ta+(1-t)b)^2} dt - \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t(1-t)}{(ta+(1-t)b)^2} dt \\
&= b^{-2} \int_0^1 t(1-t) \left(1 - t \left(1 - \frac{a}{b} \right) \right)^{-2} dt - \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} dt \\
&\quad + \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^2}{(ta+(1-t)b)^2} dt \\
&= \left[\begin{array}{l} \frac{b^{-2}}{6} {}_2F_1 \left(2, 2; 4; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{2} \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \frac{b}{3} \left(\frac{(x-a)}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right] = T_4(a, b, x). \tag{10}
\end{aligned}$$

A combination of (6)-(10) we have (5). This completes the proof.

Corollary 1. In addition to the conditions of the Theorem 3, if we choose:

1. $|f'(x)| \leq M$, for all $x \in [a, b]$, we have the following Ostrowski's type inequality

$$\left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a)M \left[\begin{array}{l} T_1(a, b, x) + T_3(a, b, x) \\ + T_2(a, b, x) + T_4(a, b, x) \end{array} \right], \quad (11)$$

2. $x = \frac{2ab}{a+b}$, we have the following midpoint type inequality for harmonically convex functions

$$\left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a) \left[\begin{array}{l} |f'(a)| (T_1(a, b, \frac{2ab}{a+b}) + T_3(a, b, \frac{2ab}{a+b})) \\ + |f'(b)| (T_2(a, b, \frac{2ab}{a+b}) + T_4(a, b, \frac{2ab}{a+b})) \end{array} \right]. \quad (12)$$

Theorem 4. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $a, b \in I$ with $a < b$ and $f' \in L$. If $|f'|^q$ is harmonically convex on $[a, b]$ for $q \geq 1$, then for all $x \in [a, b]$, we have

$$\begin{aligned} \left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a) & \left[\left(\frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right)^{1-\frac{1}{q}} \left(\begin{array}{l} |f'(a)|^q T_5(a, b, x) \\ + |f'(b)|^q T_6(a, b, x) \end{array} \right)^{\frac{1}{q}} \right. \\ & \left. + \left(\frac{1}{2} + \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 - \frac{(x-a)b}{(b-a)x} \right)^{1-\frac{1}{q}} \left(\begin{array}{l} |f'(a)|^q T_7(a, b, x) \\ + |f'(b)|^q T_8(a, b, x) \end{array} \right)^{\frac{1}{q}} \right] \end{aligned} \quad (13)$$

where

$$T_5(a, b, x) = \left[\begin{array}{l} \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{3b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^3 {}_2F_1 \left(2q, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right],$$

$$T_6(a, b, x) = \frac{1}{3b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^3 {}_2F_1 \left(2q, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right),$$

$$T_7(a, b, x) = \left[\begin{array}{l} \frac{1}{3b^{2q}} {}_2F_1 \left(2q, 1; 4; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{b^{2q}} \frac{(x-a)b}{(b-a)x} {}_2F_1 \left(2q, 1; 2; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \frac{1}{b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{3b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^3 {}_2F_1 \left(2q, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right],$$

$$T_8(a, b, x) = \left[\begin{array}{l} \frac{1}{6b^{2q}} {}_2F_1 \left(2q, 2; 4; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \frac{1}{3b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right].$$

Proof. By using Lemma 1, power mean inequality and harmonically convexity of $|f'|^q$, we have

$$\left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a) \left[\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} \left| f' \left(\frac{ab}{ta+(1-t)b} \right) \right| dt \right]$$

$$\begin{aligned}
& + \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^2} \left| f' \left(\frac{ab}{ta+(1-t)b} \right) \right| dt \\
& \leq ab(b-a) \left[\left(\int_0^{\frac{(x-a)b}{(b-a)x}} t dt \right)^{1-\frac{1}{q}} \left(\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^{2q}} [(1-t)|f'(a)|^q + t|f'(b)|^q] dt \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 (1-t) dt \right)^{1-\frac{1}{q}} \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^{2q}} [(1-t)|f'(a)|^q + t|f'(b)|^q] dt \right)^{\frac{1}{q}} \right] \\
& \leq ab(b-a) \left[\left(\frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right)^{1-\frac{1}{q}} \left(|f'(a)|^q \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t(1-t)}{(ta+(1-t)b)^{2q}} dt \right)^{\frac{1}{q}} \right. \\
& \quad \left. + |f'(b)|^q \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^2}{(ta+(1-t)b)^{2q}} dt \right)^{\frac{1}{q}} \\
& \quad + \left(\frac{1}{2} + \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 - \frac{(x-a)b}{(b-a)x} \right)^{1-\frac{1}{q}} \left(|f'(a)|^q \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{(1-t)^2}{(ta+(1-t)b)^{2q}} dt \right)^{\frac{1}{q}} \\
& \quad \left. + |f'(b)|^q \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{t(1-t)}{(ta+(1-t)b)^{2q}} dt \right)^{\frac{1}{q}}. \tag{14}
\end{aligned}$$

Calculating appearing integrals with hypergeometric functions, we have

$$\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t(1-t)}{(ta+(1-t)b)^{2q}} dt = \left[\begin{array}{l} \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{3b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^3 {}_2F_1 \left(2q, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right] = T_5(a, b, x), \tag{15}$$

$$\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^2}{(ta+(1-t)b)^{2q}} dt = \frac{1}{3b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^3 {}_2F_1 \left(2q, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) = T_6(a, b, x), \tag{16}$$

$$\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{(1-t)^2}{(ta+(1-t)b)^{2q}} dt = \left[\begin{array}{l} \frac{1}{3b^{2q}} {}_2F_1 \left(2q, 1; 4; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right) {}_2F_1 \left(2q, 1; 2; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \frac{1}{b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{3b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^3 {}_2F_1 \left(2q, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right] = T_7(a, b, x), \tag{17}$$

$$\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{t(1-t)}{(ta+(1-t)b)^{2q}} dt = \left[\begin{array}{l} \frac{1}{6b^{2q}} {}_2F_1 \left(2q, 2; 4; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \frac{1}{3b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^3 {}_2F_1 \left(2, 3; 4; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right] = T_8(a, b, x). \tag{18}$$

A combination of (14)-(18) we have (13). This completes the proof.

Corollary 2. In addition to the conditions of the Theorem 4, if we choose:

(1) $|f'(x)| \leq M$, for all $x \in [a, b]$, we have the following Ostrowski's type inequality

$$\left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a)M \left[\left(\frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right)^{1-\frac{1}{q}} (T_5(a, b, x) + T_6(a, b, x))^{\frac{1}{q}} \right. \\ \left. + \left(\frac{1}{2} + \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 - \frac{(x-a)b}{(b-a)x} \right)^{1-\frac{1}{q}} (T_7(a, b, x) + T_8(a, b, x))^{\frac{1}{q}} \right], \quad (19)$$

(2) $x = \frac{2ab}{a+b}$, we have the following midpoint type inequality for harmonically convex functions

$$\left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a) \left(\frac{1}{8} \right)^{1-\frac{1}{q}} \left[\left(|f'(a)|^q T_5\left(a, b, \frac{2ab}{a+b}\right) + |f'(b)|^q T_6\left(a, b, \frac{2ab}{a+b}\right) \right)^{\frac{1}{q}} \right. \\ \left. + \left(|f'(a)|^q T_7\left(a, b, \frac{2ab}{a+b}\right) + |f'(b)|^q T_8\left(a, b, \frac{2ab}{a+b}\right) \right)^{\frac{1}{q}} \right]. \quad (20)$$

Theorem 5. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $a, b \in I$ with $a < b$ and $f' \in L$. If $|f'|^q$ is harmonically convex on $[a, b]$ for $q \geq 1$, then for all $x \in [a, b]$, we have

$$\left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a) \left[(T_9(a, b, x))^{1-\frac{1}{q}} (|f'(a)|^q T_1(a, b, x) + |f'(b)|^q T_2(a, b, x))^{\frac{1}{q}} \right. \\ \left. + (T_{10}(a, b, x))^{1-\frac{1}{q}} (|f'(a)|^q T_3(a, b, x) + |f'(b)|^q T_4(a, b, x))^{\frac{1}{q}} \right] \quad (21)$$

where $T_1(a, b, x) - T_4(a, b, x)$ are defined as in Theorem 3 and

$$T_9(a, b, x) = \frac{1}{2} \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right),$$

$$T_{10}(a, b, x) = \begin{bmatrix} \frac{b^{-2}}{2} {}_2F_1 \left(2, 1; 3; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{(x-a)b^{-1}}{(b-a)x} {}_2F_1 \left(2, 1; 2; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \frac{1}{2} \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{bmatrix}.$$

Proof. By using Lemma 1, power mean inequality and harmonically convexity of $|f'|^q$, we have

$$\left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a) \left[\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} \left| f'\left(\frac{ab}{ta+(1-t)b}\right) \right| dt \right. \\ \left. + \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^2} \left| f'\left(\frac{ab}{ta+(1-t)b}\right) \right| dt \right] \\ \leq ab(b-a) \left[\left(\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} dt \right)^{1-\frac{1}{q}} \left(\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} [(1-t)|f'(a)|^q + t|f'(b)|^q] dt \right)^{\frac{1}{q}} \right. \\ \left. + \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^2} dt \right)^{1-\frac{1}{q}} \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^2} [(1-t)|f'(a)|^q + t|f'(b)|^q] dt \right)^{\frac{1}{q}} \right]$$

$$\begin{aligned} &\leq ab(b-a) \left[\left(\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} dt \right)^{1-\frac{1}{q}} \left(|f'(a)|^q \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t(1-t)}{(ta+(1-t)b)^{2q}} dt \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^2} dt \right)^{1-\frac{1}{q}} \left(|f'(a)|^q \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{(1-t)^2}{(ta+(1-t)b)^{2q}} dt \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(|f'(b)|^q \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{t(1-t)}{(ta+(1-t)b)^{2q}} dt \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (22)$$

Calculating appearing integrals with hypergeometric functions, we have

$$\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} dt = \frac{1}{2} \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) = T_9(a, b, x), \quad (23)$$

$$\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^2} dt = \begin{bmatrix} \frac{b^{-2}}{2} {}_2F_1 \left(2, 1; 3; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{(x-a)b^{-1}}{(b-a)x} {}_2F_1 \left(2, 1; 2; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \frac{1}{2} \left(\frac{(x-a)}{(b-a)x} \right)^2 {}_2F_1 \left(2, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{bmatrix} = T_{10}(a, b, x). \quad (24)$$

A combination of (7)-(10) and (22)-(24) we have (21). This completes the proof.

Corollary 3. In addition to the conditions of the Theorem 5, if we choose:

(1) $|f'(x)| \leq M$, for all $x \in [a, b]$, we have the following Ostrowski's type inequality

$$\begin{aligned} \left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| &\leq ab(b-a)M \left[(T_9(a, b, x))^{1-\frac{1}{q}} (T_1(a, b, x) + T_2(a, b, x))^{\frac{1}{q}} \right. \\ &\quad \left. + (T_{10}(a, b, x))^{1-\frac{1}{q}} (T_3(a, b, x) + T_4(a, b, x))^{\frac{1}{q}} \right], \end{aligned} \quad (25)$$

(2) $x = \frac{2ab}{a+b}$, we have the following midpoint type inequality for harmonically convex functions

$$\begin{aligned} &\left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \\ &\leq ab(b-a) \left[\left(T_9\left(a, b, \frac{2ab}{a+b}\right) \right)^{1-\frac{1}{q}} \left(|f'(a)|^q T_1\left(a, b, \frac{2ab}{a+b}\right) + |f'(b)|^q T_2\left(a, b, \frac{2ab}{a+b}\right) \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(T_{10}\left(a, b, \frac{2ab}{a+b}\right) \right)^{1-\frac{1}{q}} \left(|f'(a)|^q T_3\left(a, b, \frac{2ab}{a+b}\right) + |f'(b)|^q T_4\left(a, b, \frac{2ab}{a+b}\right) \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (26)$$

Theorem 6. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $a, b \in I$ with $a < b$ and $f' \in L$. If $|f'|^q$ is harmonically convex on $[a, b]$ for $q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$, then for all $x \in [a, b]$, we have

$$\begin{aligned} \left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| &\leq ab(b-a) \left[\left(\frac{1}{p+1} \left(\frac{(x-a)b}{(b-a)x} \right)^{p+1} \right)^{\frac{1}{p}} \left(|f'(a)|^q T_{11}(a, b, x) \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\frac{1}{p+1} \left(\frac{(b-x)a}{(b-a)x} \right)^{p+1} \right)^{\frac{1}{p}} \left(|f'(a)|^q T_{13}(a, b, x) \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(|f'(b)|^q T_{12}(a, b, x) \right)^{\frac{1}{q}} \right] \end{aligned} \quad (27)$$

where

$$T_{11}(a, b, x) = \left[\begin{array}{l} \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right) {}_2F_1 \left(2q, 1; 2; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right],$$

$$T_{12}(a, b, x) = \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right),$$

$$T_{13}(a, b, x) = \left[\begin{array}{l} \frac{1}{3b^{2q}} {}_2F_1 \left(2q, 1; 3; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right) {}_2F_1 \left(2q, 1; 2; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ + \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right],$$

$$T_{14}(a, b, x) = \left[\begin{array}{l} \frac{1}{2b^{2q}} {}_2F_1 \left(2q, 2; 3; \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right].$$

Proof. By using Lemma 1, Hölder inequality and harmonically convexity of $|f'|^q$, we have

$$\begin{aligned} & \left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \\ & \leq ab(b-a) \left[\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^2} \left| f' \left(\frac{ab}{ta+(1-t)b} \right) \right| dt + \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^2} \left| f' \left(\frac{ab}{ta+(1-t)b} \right) \right| dt \right] \\ & \leq ab(b-a) \left[\left(\int_0^{\frac{(x-a)b}{(b-a)x}} t^p dt \right)^{\frac{1}{p}} \left(\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{1}{(ta+(1-t)b)^{2q}} [(1-t)|f'(a)|^q + t|f'(b)|^q] dt \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 (1-t)^p dt \right)^{\frac{1}{p}} \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^{2q}} [(1-t)|f'(a)|^q + t|f'(b)|^q] dt \right)^{\frac{1}{q}} \right] \\ & \leq ab(b-a) \left[\left(\frac{1}{p+1} \left(\frac{(x-a)b}{(b-a)x} \right)^{p+1} \right)^{\frac{1}{p}} \left(\begin{array}{l} |f'(a)|^q \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{1-t}{(ta+(1-t)b)^{2q}} dt \\ + |f'(b)|^q \int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta+(1-t)b)^{2q}} dt \end{array} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\frac{1}{p+1} \left(\frac{(b-x)a}{(b-a)x} \right)^{p+1} \right)^{\frac{1}{p}} \left(\begin{array}{l} |f'(a)|^q \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta+(1-t)b)^{2q}} dt \\ + |f'(b)|^q \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{t}{(ta+(1-t)b)^{2q}} dt \end{array} \right)^{\frac{1}{q}} \right]. \end{aligned} \tag{28}$$

Calculating appearing integrals with hypergeometric functions, we have

$$\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{1-t}{(ta+(1-t)b)^{2q}} dt = \left[\begin{array}{l} \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right) {}_2F_1 \left(2q, 1; 2; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \\ - \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) \end{array} \right] = T_{11}(a, b, x), \tag{29}$$

$$\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta + (1-t)b)^{2q}} dt = \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} (1 - \frac{a}{b}) \right) = T_{12}(a, b, x), \quad (30)$$

$$\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta + (1-t)b)^{2q}} dt = \begin{bmatrix} \frac{1}{3b^{2q}} {}_2F_1 \left(2q, 1; 3; \left(1 - \frac{a}{b} \right) \right) \\ -\frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right) {}_2F_1 \left(2q, 1; 2; \frac{(x-a)b}{(b-a)x} (1 - \frac{a}{b}) \right) \\ + \frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} (1 - \frac{a}{b}) \right) \end{bmatrix} = T_{13}(a, b, x), \quad (31)$$

$$\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{t}{(ta + (1-t)b)^{2q}} dt = \begin{bmatrix} \frac{1}{2b^{2q}} {}_2F_1 \left(2q, 2; 3; \left(1 - \frac{a}{b} \right) \right) \\ -\frac{1}{2b^{2q}} \left(\frac{(x-a)b}{(b-a)x} \right)^2 {}_2F_1 \left(2q, 2; 3; \frac{(x-a)b}{(b-a)x} (1 - \frac{a}{b}) \right) \end{bmatrix} = T_{14}(a, b, x). \quad (32)$$

A combination of (28)-(32) we have (27). This completes the proof.

Corollary 4. In addition to the conditions of the Theorem 6, if we choose:

(1) $|f'(x)| \leq M$, for all $x \in [a, b]$, we have the following Ostrowski's type inequality

$$\begin{aligned} \left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| &\leq ab(b-a)M \left[\left(\frac{1}{p+1} \left(\frac{(x-a)b}{(b-a)x} \right)^{p+1} \right)^{\frac{1}{p}} (T_{11}(a, b, x) + T_{12}(a, b, x))^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\frac{1}{p+1} \left(\frac{(b-x)a}{(b-a)x} \right)^{p+1} \right)^{\frac{1}{p}} (T_{13}(a, b, x) + T_{14}(a, b, x))^{\frac{1}{q}} \right], \end{aligned} \quad (33)$$

(2) $x = \frac{2ab}{a+b}$, we have the following midpoint type inequality for harmonically convex functions

$$\begin{aligned} \left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| &\leq ab(b-a) \left[\left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(|f'(a)|^q T_{11}(a, b, \frac{2ab}{a+b}) + |f'(b)|^q T_{12}(a, b, \frac{2ab}{a+b}) \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(|f'(a)|^q T_{13}(a, b, \frac{2ab}{a+b}) + |f'(b)|^q T_{14}(a, b, \frac{2ab}{a+b}) \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (34)$$

Theorem 7. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $a, b \in I$ with $a < b$ and $f' \in L$. If $|f'|^q$ is harmonically convex on $[a, b]$ for $q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$, then for all $x \in [a, b]$, we have

$$\begin{aligned} \left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| &\leq ab(b-a) \left[(T_{15}(a, b, x))^{\frac{1}{p}} \left(|f'(a)|^q \left[\frac{(x-a)b}{(b-a)x} - \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right] \right)^{\frac{1}{q}} \right. \\ &\quad \left. + |f'(b)|^q \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right)^{\frac{1}{q}} \\ &\quad + (T_{16}(a, b, x))^{\frac{1}{p}} \left(|f'(a)|^q \left[\frac{1}{2} - \frac{(x-a)b}{(b-a)x} + \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right] \right)^{\frac{1}{q}} \\ &\quad \left. + |f'(b)|^q \left[\frac{1}{2} - \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right] \right)^{\frac{1}{q}} \right] \end{aligned} \quad (35)$$

where

$$\begin{aligned} T_{15}(a, b, x) &= \frac{1}{2b^{2p}} \left(\frac{(x-a)b}{(b-a)x} \right)^{p+1} {}_2F_1 \left(2p, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right), \\ T_{16}(a, b, x) &= \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{(1-t)^p}{(ta + (1-t)b)^{2p}} dt. \end{aligned}$$

Proof. By using Lemma 1, Hölder inequality and harmonically convexity of $|f'|^q$, we have

$$\begin{aligned} \left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| &\leq ab(b-a) \left[\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t}{(ta + (1-t)b)^2} \left| f' \left(\frac{ab}{ta + (1-t)b} \right) \right| dt \right. \\ &\quad \left. + \int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{1-t}{(ta + (1-t)b)^2} \left| f' \left(\frac{ab}{ta + (1-t)b} \right) \right| dt \right] \\ &\leq ab(b-a) \left[\left(\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^p}{(ta + (1-t)b)^{2p}} dt \right)^{\frac{1}{p}} \left(\int_0^{\frac{(x-a)b}{(b-a)x}} [(1-t)|f'(a)|^q + t|f'(b)|^q] dt \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{(1-t)^p}{(ta + (1-t)b)^{2p}} dt \right)^{\frac{1}{p}} \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 [(1-t)|f'(a)|^q + t|f'(b)|^q] dt \right)^{\frac{1}{q}} \right] \\ &\leq ab(b-a) \left[\left(\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^p}{(ta + (1-t)b)^{2p}} dt \right)^{\frac{1}{p}} \left(\begin{array}{l} |f'(a)|^q \left[\frac{(x-a)b}{(b-a)x} - \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right] \\ + |f'(b)|^q \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \end{array} \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{(1-t)^p}{(ta + (1-t)b)^{2p}} dt \right)^{\frac{1}{p}} \left(\begin{array}{l} |f'(a)|^q \left[\frac{1}{2} - \frac{(x-a)b}{(b-a)x} + \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right] \\ + |f'(b)|^q \left[\frac{1}{2} - \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right] \end{array} \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (36)$$

Since the appearing integrals are as the following, we have

$$\int_0^{\frac{(x-a)b}{(b-a)x}} \frac{t^p}{(ta + (1-t)b)^{2p}} dt = \frac{1}{2b^{2p}} \left(\frac{(x-a)b}{(b-a)x} \right)^{p+1} {}_2F_1 \left(2p, 2; 3; \frac{(x-a)b}{(b-a)x} \left(1 - \frac{a}{b} \right) \right) = T_{15}(a, b, x), \quad (37)$$

$$\int_{\frac{(x-a)b}{(b-a)x}}^1 \frac{(1-t)^p}{(ta + (1-t)b)^{2p}} dt = T_{16}(a, b, x). \quad (38)$$

A combination of (36)-(38) we have (35). This completes the proof.

Corollary 5. In addition to the conditions of the Theorem 5, if we choose:

(1) $|f'(x)| \leq M$, for all $x \in [a, b]$, we have the following Ostrowski's type inequality

$$\begin{aligned} \left| f(x) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| &\leq ab(b-a)M \left[\left(T_{15}(a, b, x) \right)^{\frac{1}{p}} \left(\left[\frac{(x-a)b}{(b-a)x} - \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right] + \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(T_{16}(a, b, x) \right)^{\frac{1}{p}} \left(\left[\frac{1}{2} - \frac{(x-a)b}{(b-a)x} + \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right] + \left[\frac{1}{2} - \frac{1}{2} \left(\frac{(x-a)b}{(b-a)x} \right)^2 \right] \right)^{\frac{1}{q}} \right], \end{aligned} \quad (39)$$

(2) $x = \frac{2ab}{a+b}$, we have the following midpoint type inequality for harmonically convex functions

$$\left| f\left(\frac{2ab}{a+b}\right) - \frac{ab}{b-a} \int_a^b \frac{f(u)}{u^2} du \right| \leq ab(b-a) \left[\left(T_{15} \left(a, b, \frac{2ab}{a+b} \right) \right)^{\frac{1}{p}} \left(\frac{3|f'(a)|^q}{8} + \frac{|f'(b)|^q}{8} \right)^{\frac{1}{q}} \right. \\ \left. + \left(T_{16} \left(a, b, \frac{2ab}{a+b} \right) \right)^{\frac{1}{p}} \left(\frac{|f'(a)|^q}{8} + \frac{3|f'(b)|^q}{8} \right)^{\frac{1}{q}} \right]. \quad (40)$$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors have contributed to all parts of the article. All authors read and approved the final manuscript.

References

- [1] M. Alomari, M. Darus, Some Ostrowski's type inequalities for convex functions with applications, *RGMIA Res.Rep. Collect.*, 13(1) (2010), Article 3, 1-14.
- [2] S. S. Dragomir, T. M. Rassias, *Ostrowski type inequalities and applications in numerical integration*, Kluwer Academic Publishers, 2002.
- [3] İ. İşcan, Hermite-Hadamard type inequalities for harmonically convex functions, *Hacet. J. Math. Stat.*, 43 (6) (2014), 935-942.
- [4] İ. İşcan, Ostrowski type inequalities for harmonically s -convex functions, *Konuralp Jurnal of Mathematics*, Volume 3, No 1 (2015) 63-74.
- [5] İ. İşcan, S. Numan, Ostrowski type inequalities for harmonically quasi-convex functions, *Elect. J. Math. Anal. App.*, 2(2) (2014) 189-198.
- [6] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, 2006
- [7] A. Ostrowski, Über die Absolutabweichung einer differentiablen Funktionen von ihren Integralmittelwert. *Comment. Math. Hel.*, 10 (1938), 226–227.
- [8] M. E. Özdemir, Ç. Yıldız, New Ostrowski Type Inequalities for Geometrically Convex Functions, *Int. J. Modern Math. Sci.*, 8(1) (2013) 27-35.
- [9] E. Set, M. E. Özdemir, M. Z. Sarıkaya, New inequalities of Ostrowski's type for s -convex functions in the second sense with applications, *Facta Uni. Ser. Math. Inform.* 27(1) (2012) 67-82.